



Upper bound Estimates of Fourth Order Hankel and Toeplitz Determinants for Certain Analytic Functions Connected with Three Leaf Function

Rudrani Rangu^{*1} , Bharavi Sharma Rayaprolu² and Sambasiva Rao Siginam³

¹ Department of Mathematics, Government Degree College (Autonomous), Narsampet, Warangal 506132, Telangana, India

² Department of Mathematics, Kakatiya University, Warangal 506009, Telangana, India

³ Department of Humanities and Sciences, SVS Group of Institutions, Warangal 506015, Telangana, India

*Corresponding author: rangurudrani1981@gmail.com

Received: October 20, 2024

Revised: January 17, 2025

Accepted: February 6, 2025

Abstract. The purpose of this paper is to compute upper bounds of Hankel and Toeplitz determinants up to fourth order for some normalized univalent functions defined on the open unit disk in the complex plane associated with three leaf function. Suitable examples are provided in support of proven results.

Keywords. Univalent functions, Starlike functions, Coefficient inequalities, Hankel determinants, Toeplitz determinants, Three Leaf domain

Mathematics Subject Classification (2020). 30C45, 30C80

Copyright © 2025 Rudrani Rangu, Bharavi Sharma Rayaprolu and Sambasiva Rao Siginam. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

1. Introduction

1.1 Preliminaries

Let \mathcal{A} represent the family of analytic functions f with normalization $f(0) = 0$ and $f'(0) = 1$ defined on the open unit disk \mathbb{D} in the complex plane \mathbb{C} . The Taylor series expansion of $f \in \mathcal{A}$ is of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad \text{for all } z \in \mathbb{D}, \text{ where } a_n = \frac{f^{(n)}(0)}{n!}. \quad (1.1)$$

The set of functions $f \in \mathcal{A}$ that are univalent is represented by \mathcal{S} . The set \mathcal{B} of analytic functions $w(z)$ defined on \mathbb{D} with the conditions $w(0) = 0$ and $|w(z)| < 1$, for all $z \in \mathbb{D}$ is refereed as the class of Schwarz functions (Goodman [7]).

A function $f \in \mathcal{A}$ is said to be subordinate to $g \in \mathcal{A}$ if there exists $w \in \mathcal{B}$ such that $f(z) = g(w(z))$, for all $z \in \mathbb{D}$. We denote it by $f < g$. In particular if g is univalent, then f is subordinate to g if, and only if, $g(0) = f(0)$ and $f(\mathbb{D}) \subset g(\mathbb{D})$. For more information regarding univalent function theory, we refer to Pommerenke [22].

The set of starlike and convex functions respectively \mathcal{S}^* , \mathcal{C} which are subclasses of \mathcal{S} and further extended by Ma and Minda [15] by utilizing the concept of subordination to $\mathcal{S}^*(\varphi)$ and $\mathcal{C}(\varphi)$ where $\varphi \in \mathcal{A}$ satisfying conditions $\Re\{\varphi(z)\} > 0$, $\varphi'(0) > 0$, $\varphi(\mathbb{D})$ is symmetric with respect to real axis and starlike with respect to $\varphi(0) = 1$. Some Ma-Minda type subclasses of \mathcal{S} were studied by choosing specific function φ in the recent past (Arif *et al.* [3], Gandhi [6], and Sharma *et al.* [27]).

The q th Hankel determinant of index $n \geq 1$ for a function $f \in \mathcal{A}$, with a series expansion given by (1.1), is denoted as $H_{q,n}(f)$ (or simply $H_q(n)$). It is defined as follows:

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \dots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \dots & a_{n+q} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n+q-1} & a_{n+q} & \dots & a_{n+2q-2} \end{vmatrix}, \quad (1.2)$$

where $q \geq 2$ and $a_1 = 1$ (refer to Pommerenke [20, 21]).

Similarly, the q th symmetric Toeplitz determinant $T_q(n)$ for a function $f \in \mathcal{A}$ is defined as follows:

$$T_q(n) = \begin{vmatrix} a_n & a_{n+1} & \dots & a_{n+q-1} \\ a_{n+1} & a_n & \dots & a_{n+q-2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n+q-1} & a_{n+q-2} & \dots & a_n \end{vmatrix}, \quad (1.3)$$

where $q \geq 2$, $n \geq 1$, and $a_1 = 1$ (refer to Ali *et al.* [1]).

In 1960, Lawrence Zalcman conjecture that the coefficients of $f \in \mathcal{S}$ with series representation (1.1) satisfy the inequality (see, Libera and Ziotkiewicz [14])

$$|a_n^2 - a_{2n-1}| \leq (n-1)^2, \quad \text{for } n \geq 2.$$

Ma [16] proposed generalized Zalcman conjecture for $f \in \mathcal{S}$ of the form (1.1) that

$$|a_n a_m - a_{n+m-1}| \leq (n-1)(m-1), \quad \text{for } n, m \geq 2$$

and proved this conjecture is true for starlike functions and univalent function with real coefficients.

1.2 Literature Review Concerning Hankel and Toeplitz Determinants in Geometric Function Theory

The pioneering works of Pommerenke [20, 21], Hayman [8], Babalola [5], Zaprawa [36], Kowalezyk *et al.* [13] on second and third Hankel determinants inspired Arif *et al.* [2] to estimate an upper bound of $|H_{4,1}(f)|$, for $f \in \mathcal{R}$. Subsequently, Srivastava *et al.* [31], Khan *et*

al. [10,11], and Yakaiah *et al.* [34] computed upper bound for $|H_{4,1}(f)|$ for f being the member of subclasses of \mathcal{S} subordinate to cardioid, sine, modified sigmoid and cosine functions, respectively.

Ali *et al.* [1] studied the Toeplitz determinant $T_q(n)$ for the class \mathcal{S} and certain of its subclasses. Zhang and Tang [37] studied the upper bounds of the fourth Toeplitz determinant for the class

$$\mathcal{S}^*(\sin) = \left\{ f \in \mathcal{S} : \frac{zf'(z)}{f(z)} < 1 + \sin(z) \right\}.$$

Vijayalakshmi *et al.* [32] studied symmetric Toeplitz determinants for classes defined by post quantum operators subordinated to the limaçon function. Mandal *et al.* [17] investigated Toeplitz determinants of logarithmic coefficients of inverse functions for certain classes of univalent function. For similar type of studies concerning Toeplitz determinant $|T_q(n)|$ for starlike, convex and bounded boundary rotation functions, we refer to Ali *et al.* [1] and Radhika *et al.* [23].

We now provide some more details of recent research done in this direction.

- (1) Kaur and Singh [9] proved $|H_{4,1}(f)| \leq \frac{4027899}{896000000}$ for the members of

$$\mathcal{R}_1 = \{f \in \mathcal{A} : \Re\{f'(z) + zf''(z)\} > 0\}.$$

- (2) Yakaiah and Sharma [33] estimated $|H_{4,1}(f)| \leq 0.136765285$, $|T_4(1)(f)| \leq 0.0069$ and $|T_4(2)(f)| \leq 0.00009$ for the members of

$$\mathcal{R}_1(\cos z) = \{f \in \mathcal{A} : f'(z) + zf''(z) < \cos z\}.$$

- (3) Koride *et al.* [12] computed $|H_{4,1}(f)| \leq 1.540436$, $|T_4(1)(f)| \leq 2.363923$ and $|T_4(2)(f)| \leq 0.140203$ for the members of

$$\mathcal{R}_1(1 + \sin z) = \{f \in \mathcal{A} : f'(z) + zf''(z) < 1 + \sin z\}.$$

- (4) Yakaiah *et al.* [34] computed $|H_{4,1}(f)| \leq 1.24199978975$, $|T_4(1)(f)| \leq 2.4375$ and $|T_4(2)(f)| \leq 0.674461806$ for the members of

$$\mathcal{R}_1(1 + \tanh z) = \{f \in \mathcal{A} : f'(z) + zf''(z) < 1 + \tanh z\}.$$

1.3 Identification of Research Problem

The Ma-Minda type function $\varphi_{3L}(z) = 1 + (4/5)z + (1/5)z^4$ maps \mathbb{D} onto three leaf shaped domain and the class

$$\mathcal{S}_{3L}^* = \left\{ f \in \mathcal{S} : \frac{zf'(z)}{f(z)} < \varphi_{3L}(z) \right\},$$

have been studied by Gandhi *et al.* [6], Shi *et al.* [28, 30]. Arif *et al.* [4] studied the bounds for third order Hankel determinant for two subfamilies of starlike and bounded turning functions associated with a three-leaf shaped domain. Raza *et al.* [26] estimated the second and third Hankel determinants for starlike and convex functions associated to three leaf function. Murugusundaramoorthy *et al.* [18] developed a new class of bi-starlike functions subordinate to a three leaf function induced by multiplicative calculus. Murugusundaramoorthy *et al.* [19] computed upper bound of $|H_3(1)|$, for the members of \mathcal{R}_{3L} . Shi *et al.* [29] studied the bounds of second Hankel determinants and with a logarithmic coefficient as entry for the class of bounded turning functions connected with a three-leaf shaped domain.

Motivated by the earlier mentioned research works, in this paper, we compute the upper bounds of fourth Hankel and Toeplitz determinants for the members of

$$\mathcal{R}_1(\varphi_{3L}) = \left\{ f \in \mathcal{S} : f'(z) + zf''(z) < \varphi_{3L}(z) = 1 + \frac{4}{5}z + \frac{1}{5}z^4, \text{ for all } z \in \mathbb{D} \right\},$$

were computed in this paper.

2. A Set of Essential Lemmas

The class of analytic functions $p(z)$ defined on the unit disk \mathbb{D} with $p(0) = 1$ and $\Re\{p(z)\} > 0$ is called the class of functions with positive real part, denoted by \mathcal{P} . For $p \in \mathcal{P}$, the Taylor series expansion of $p \in \mathcal{P}$ is

$$p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n, \quad \text{for } z \in \mathbb{D}. \quad (2.1)$$

Unless otherwise stated throughout this paper, we assume the series representation of $p \in \mathcal{P}$ is of the form (2.1).

Lemma 2.1 ([22]). *Let $p \in \mathcal{P}$. Then $|c_n| \leq 2$ for any positive integer n . The inequality is sharp for $p(z) = \frac{1+z}{1-z}$.*

Lemma 2.2 ([15]). *Let $p \in \mathcal{P}$ and $\eta \in \mathbb{C}$. Then $|c_2 - \eta c_1^2| \leq 2 \max\{1, |2\eta - 1|\}$. The inequality is sharp for $p(z) = \frac{1+z}{1-z}$ and $p(z) = \frac{1+z^2}{1-z^2}$.*

Lemma 2.3 ([3]). *Let $p \in \mathcal{P}$. Then for any real numbers A, B and C ,*

$$|Ac_1^3 - Bc_1c_2 + Cc_3| \leq 2|A| + 2|B - 2A| + 2|A - B + C|.$$

Lemma 2.4 ([25]). *Let $p \in \mathcal{P}$. Then for all $n, m \in \mathbb{N}$,*

$$|\eta c_n c_m - c_{n+m}| = \begin{cases} 2, & \text{if } 0 \leq \eta \leq 1, \\ 2|2\eta - 1|, & \text{otherwise.} \end{cases}$$

This inequality is sharp.

Lemma 2.5 ([25]). *Let $p \in \mathcal{P}$ and l, m, n and r be real numbers and if the inequalities $0 < m < 1$, $0 < r < 1$,*

$$8r(1-r)((mn - 2l)^2 + (m(r+m) - n)^2) + m(1-m)(n - 2rm)^2 \leq 4m^2(1-m)^2r(1-r) \quad (2.2)$$

hold, then

$$\left| lc_1^4 + rc_2^2 + 2mc_1c_3 - \frac{3n}{2}c_1^2c_2 - c_4 \right| \leq 2. \quad (2.3)$$

Lemma 2.6 ([14]). *If the function $p \in \mathcal{P}$, then*

$$2c_2 = c_1^2 + x(4 - c_1^2), \quad (2.4)$$

$$4c_3 = c_1^3 + 2(4 - c_1^2)c_1x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)z, \quad (2.5)$$

for some x, z with $|x| \leq 1$ and $|z| \leq 1$.

3. Main Results

Let $f \in \mathcal{R}_1(\varphi_{3L})$ be of the form (1.1). Then there exists $w \in \mathcal{B}$ such that

$$f'(z) + zf''(z) = \varphi_{3L}(w(z)) = 1 + \frac{4}{5}w(z) + \frac{1}{5}w(z)^4, \quad \text{for all } z \in \mathbb{D}. \quad (3.1)$$

If we take $p(z) = \frac{1+w(z)}{1-w(z)}$, for all $z \in \mathbb{D}$ then $p \in \mathcal{P}$ and $w(z) = \frac{p(z)-1}{p(z)+1}$ so that

$$f'(z) + zf''(z) = 1 + \frac{4}{5} \left(\frac{p(z)-1}{p(z)+1} \right) + \frac{1}{5} \left(\frac{p(z)-1}{p(z)+1} \right)^4, \quad \text{for all } z \in \mathbb{D}. \quad (3.2)$$

Substituting (1.1) and (2.1) in (3.2) followed by comparing like coefficients on both sides of (3.2), we obtain

$$a_2 = \frac{c_1}{10}, \quad (3.3)$$

$$a_3 = \frac{2}{45} \left(c_2 - \frac{c_1^2}{2} \right), \quad (3.4)$$

$$a_4 = \frac{1}{40} \left(\frac{1}{4}c_1^3 - c_1c_2 + c_3 \right), \quad (3.5)$$

$$a_5 = \frac{2}{125} \left(c_4 - c_1c_3 - \frac{1}{2}c_2^2 + \frac{3}{4}c_1^2c_2 - \frac{3}{32}c_1^4 \right), \quad (3.6)$$

$$a_6 = \frac{2}{180} \left(c_5 - c_1c_4 - c_2c_3 + \frac{3}{4}c_1^2c_3 + \frac{3}{4}c_1c_2^2 - \frac{3}{8}c_1^3c_2 \right), \quad (3.7)$$

$$a_7 = \frac{2}{245} \left(c_6 - c_1c_5 - c_2c_4 - \frac{1}{2}c_3^2 + \frac{3}{4}c_1^2c_4 + \frac{3}{2}c_1c_2c_3 + \frac{1}{4}c_2^3 - \frac{3}{8}c_1^3c_3 - \frac{9}{16}c_1^2c_2^2 + \frac{3}{64}c_1^6 \right). \quad (3.8)$$

Example 3.1. By taking the Schwarz functions $w(z) = z$, $w(z) = z^2$, $w(z) = z^3$ and $w(z) = z^4$ in (3.1) followed by integrating on both sides and utilizing the fact $f(0) = 0$, $f'(0) = 1$, we get

- (1) $f_1(z) = z + \frac{1}{5}z^2 + \frac{1}{125}z^5$,
- (2) $f_2(z) = z + \frac{4}{45}z^3 + \frac{1}{405}z^9$,
- (3) $f_3(z) = z + \frac{1}{20}z^4 + \frac{1}{845}z^{13}$,
- (4) $f_4(z) = z + \frac{4}{125}z^5 + \frac{1}{1445}z^{17}$.

We can observe that all the above functions are in $\mathcal{R}_1(\varphi_{3L})$.

We now estimate initial coefficient bounds for the functions in $\mathcal{R}_1(\varphi_{3L})$.

Theorem 3.1. If $f \in \mathcal{R}_1(\varphi_{3L})$ is given by (1.1). Then $|a_2| \leq \frac{1}{5}$, $|a_3| \leq \frac{4}{45}$, $|a_4| \leq \frac{1}{20}$, $|a_5| \leq \frac{4}{125}$, $|a_6| \leq \frac{1}{10}$ and $|a_7| \leq \frac{38}{245}$. The functions f_1, f_2, f_3, f_4 as in Example 3.1 are extremal functions for first four inequalities, respectively.

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). By applying Lemmas 2.1, 2.2 and 2.3 to (3.3), (3.4) and (3.5) respectively, we obtain

$$|a_2| = \frac{|c_1|}{10} \leq \frac{1}{5},$$

$$|a_3| = \left| \frac{2}{45} \left(c_2 - \frac{c_1^2}{2} \right) \right| \leq \frac{4}{45},$$

$$|a_4| = \frac{1}{160} |c_1^3 - 4c_1c_2 + 4c_3| \leq \frac{2(1+2+1)}{160} = \frac{1}{20}.$$

Now,

$$\begin{aligned} |a_5| &= \frac{2}{125} \left| c_4 - c_1c_3 - \frac{1}{2}c_2^2 + \frac{3}{4}c_1^2c_2 - \frac{3}{32}c_1^4 \right| \\ &= \frac{2}{125} \left| lc_1^4 + rc_2^2 + 2mc_1c_3 - \frac{3n}{2}c_1^2c_2 - c_4 \right|, \end{aligned}$$

where $l = \frac{3}{32}$, $r = \frac{1}{2}$, $m = \frac{1}{2}$ and $n = \frac{1}{2}$.

These values of l, m, n and r satisfy the hypothesis of Lemma 2.5 and hence $|a_5| \leq \frac{4}{125}$.

By Lemmas 2.1, 2.4 and 2.5, we have $|c_1| \leq 2$, $|c_2| \leq 2$, $|c_5 - c_2c_3| \leq 2$ and $|\frac{3}{8}c_2^2 + \frac{3}{4}c_1c_3 - \frac{3}{8}c_1^2c_2 - c_4| \leq 2$. Consequently, we obtain the following after a simple computation

$$\begin{aligned} |a_6| &= \frac{1}{90} \left| c_5 - c_1c_4 - c_2c_3 + \frac{3}{4}c_1^2c_3 + \frac{3}{4}c_1c_2^2 - \frac{3}{8}c_1^3c_2 \right| \\ &\leq \frac{1}{90} \left(|c_5 - c_2c_3| + |c_1| \left| \frac{3}{8}c_2^2 + \frac{3}{4}c_1c_3 - \frac{3}{8}c_1^2c_2 - c_4 \right| + \frac{3}{8}|c_1||c_2|^2 \right) \\ &\leq \frac{1}{10}. \end{aligned}$$

Further, in view of Lemma 2.3, we have $|\frac{3}{8}c_1^3 + c_1c_2 - \frac{1}{2}c_3| \leq \frac{3}{2}$ and in view of Lemma 2.4, we have $|c_6 - c_1c_5| \leq 2$, $|c_4 - \frac{1}{2}c_1c_3| \leq 2$, $|c_2 - \frac{9}{4}c_1^2| \leq 7$ and using the fact $|c_n| \leq 2$ for $n \geq 1$, we obtain

$$\begin{aligned} |a_7| &= \frac{2}{245} \left| c_6 - c_1c_5 - c_2c_4 - \frac{1}{2}c_3^2 + \frac{3}{4}c_1^2c_4 + \frac{3}{2}c_1c_2c_3 + \frac{1}{4}c_2^3 - \frac{3}{8}c_1^3c_3 - \frac{9}{16}c_1^2c_2^2 + \frac{3}{64}c_1^6 \right| \\ &\leq \frac{2}{245} \left(|c_6 - c_1c_5| + \frac{|c_2|^2}{4} \left| c_2 - \frac{9}{4}c_1^2 \right| + |c_2| \left| c_4 - \frac{1}{2}c_1c_3 \right| + |c_3| \left| -\frac{3}{8}c_1^3 + c_1c_2 - \frac{1}{2}c_3 \right| + \frac{3}{64}|c_1|^6 \right) \\ &\leq \frac{38}{245}. \end{aligned}$$

We now obtain upper bound for Fekete-Szegő functional of the class $\mathcal{R}_1(\varphi_{3L})$.

Theorem 3.2. If $f \in \mathcal{R}_1(\varphi_{3L})$ is given by (1.1) and $H_{2,1}^\eta(f) = a_3 - \eta a_2^2$ for $\eta \in \mathbb{C}$, then

$$|H_{2,1}^\eta(f)| \leq \frac{4}{45} \max \left\{ 1, \frac{9}{20}|\eta| \right\} \quad (3.9)$$

and this inequality is sharp.

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$. Then in view of Lemma 2.2, we obtain

$$\begin{aligned} |H_{2,1}^\eta(f)| &= |a_3 - \eta a_2^2| \\ &= \frac{2}{45} \left| c_2 - \left(\frac{20+9\eta}{40} \right) c_1^2 \right| \\ &\leq \frac{4}{45} \max \left\{ 1, \left| 2 \left(\frac{20+9\eta}{40} \right) - 1 \right| \right\} \\ &= \frac{4}{45} \max \left\{ 1, \frac{9}{20}|\eta| \right\}. \end{aligned}$$

Sharpness is evident from the fact that if $|\eta| \leq \frac{20}{9}$, then $|H_{2,1}^\eta(f_2)| = \frac{4}{45}$ and if $|\eta| > \frac{20}{9}$, then $|H_{2,1}^\eta(f_1)| = \frac{|\eta|}{25}$, where f_1 and f_2 are as in Example 3.1. \square

We now estimate an upper bound for second Hankel determinants for the class $\mathcal{R}_1(\varphi_{3L})$.

Theorem 3.3. *If $f \in \mathcal{R}_1(\varphi_{3L})$ is given by (1.1), then $|H_{2,2}(f)| = |a_2a_4 - a_3^2| \leq \frac{16}{2025}$. Sharpness is obtained for the function f_2 as in Example 3.1.*

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). By utilizing (3.3), (3.4) and (3.5), we have

$$\begin{aligned} |H_{2,2}(f)| &= \frac{1}{25} \left| \frac{c_1}{5184} (17c_1^3 - 68c_1c_2 + 324c_3) - \frac{4}{81} c_2^2 \right| \\ &= \frac{1}{129600} |324c_1c_3 - 68c_1^2c_2 + 17c_1^4 - 256c_2^2|. \end{aligned}$$

Applying Lemma 2.6, we obtain

$$\begin{aligned} |H_{2,2}(f)| &= \frac{1}{129600} |(81 - 34 + 17 - 64)c_1^4 + (162 - 34 - 128)c_1^2x(4 - c_1^2) \\ &\quad - 81c_1^2x^2(4 - c_1^2) - 64x^2(4 - c_1^2)^2 + 162(4 - c_1^2)c_1(1 - |x|^2)z| \\ &= \frac{1}{129600} |-81c_1^2x^2(4 - c_1^2) - 64x^2(4 - c_1^2)^2 + 162(4 - c_1^2)c_1(1 - |x|^2)z| \\ &\leq \frac{1}{129600} (81c^2t^2(4 - c^2) + 64t^2(4 - c^2)^2 + 162(4 - c^2)c(1 - t^2)z) \\ &:= F(c, t), \end{aligned}$$

where $c = |c_1| \in [0, 2]$, $t = |x| \in [0, 1]$ and $0 \leq z \leq 1$. Thus,

$$\begin{aligned} \frac{\partial F}{\partial t} &= \frac{1}{64800} (4 - c^2)(17c^2 - 162c + 256)t \\ &= \frac{1}{64800} (4 - c^2)(2 - c)(128 - 17c)t \\ &\geq 0. \end{aligned}$$

Therefore, $F(c, t)$ is an increasing function in variable t and hence

$$\begin{aligned} |H_{2,2}(f)| &\leq F(c, 1) \\ &= \frac{1}{129600} (81c^2(4 - c^2) + 64(4 - c^2)^2) \\ &:= \psi(c). \end{aligned}$$

It is clear that $\psi'(c) = 0$ implies $c = 0$ and $c = \sqrt{\frac{-376}{68}}$. Since $0 \leq c \leq 2$, we take $c = 0$. Further, $\psi''(0) = -376 < 0$. Therefore, ψ attains its maximum at $c = 0$. Hence,

$$\begin{aligned} |H_{2,2}(f)| &\leq \psi(0) \\ &= \frac{1024}{129600} \\ &= \frac{16}{2025}. \end{aligned}$$

□

Theorem 3.4. *If $f \in \mathcal{R}_1(\varphi_{3L})$, then $|H_{2,3}(f)| \leq \frac{3373}{360000}$.*

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). By using (3.4), (3.5) and (3.6), we obtain

$$|H_{2,3}(f)| = \frac{1}{5760000} |4096c_2c_4 - 2048c_2^3 + 3014c_1c_2c_3 + 496c_1^2c_2^2 - 120c_1^4c_2 - 2048c_1^2c_4|$$

$$\begin{aligned}
& + 248c_1^3c_3 - 33c_1^6 + 248c_1^3c_3 - 33c_1^6 - 3600c_3^2| \\
& \leq \frac{1}{5760000} \left(4096|c_2| \left| \frac{15}{512}c_1^4 + \frac{1}{2}c_2^2 + \frac{293}{1024}c_1c_3 - \frac{31}{256}c_1^2c_2 - c_4 \right| + 3600|c_3| |c_3 - c_1c_2| \right. \\
& \quad \left. + 2048|c_1|^2 \left| c_4 - \frac{31}{256}c_1c_3 \right| + 586|c_1||c_2||c_3| + 33|c_1|^6 \right).
\end{aligned}$$

An application of Lemma 2.5 shows that

$$\left| \frac{15}{512}c_1^4 + \frac{1}{2}c_2^2 + \frac{293}{1024}c_1c_3 - \frac{31}{256}c_1^2c_2 - c_4 \right| \leq 2.$$

Further, it is clear to see that an application of Lemma 2.4 yields $|c_3 - c_1c_2| \leq 2$, $|c_4 - \frac{31}{256}c_1c_3| \leq 2$. In view of these inequalities along with $|c_n| \leq 2$ for all $n \geq 1$, we finally obtain

$$\begin{aligned}
|H_{2,3}(f)| & \leq \frac{3373}{360000} \\
& \approx 0.009369444.
\end{aligned}$$

□

3.1 Bounds of Zalcman Functionals, $|H_{3,1}(f)|$, $|H_{3,2}(f)|$ for the class $\mathcal{R}_1(\varphi_{3L})$

Theorem 3.5. If $f \in \mathcal{R}_1(\varphi_{3L})$, then $|a_4 - a_2a_3| \leq \frac{1}{20}$. The sharpness is obtained for

$$f_3(z) = z + \frac{1}{20}z^4 + \frac{z^{13}}{845} \in \mathcal{R}_1(\varphi_{3L}).$$

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). Utilizing (3.3), (3.4) and (3.5) and applying Lemma 2.3, we obtain

$$\begin{aligned}
|a_4 - a_2a_3| & = \frac{1}{7200} |77c_1^3 - 212c_1c_2 + 180c_3| \\
& \leq \frac{2}{7200} (77 + |212 - 154| + |77 - 212 + 180|) \\
& = \frac{1}{20}.
\end{aligned}$$

□

Theorem 3.6. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). Then $|a_3 - a_2^2| \leq \frac{4}{45}$, $|a_5 - a_3^2| \leq \frac{4}{125}$ and $|a_7 - a_4^2| \leq \frac{3089}{19600}$. Further, $f_2(z) = z + \frac{4}{45}z^3 + \frac{1}{405}z^9$, $f_4(z) = z + \frac{4}{125}z^5 + \frac{1}{1445}z^{17}$ as given in Example 3.1 are extremal functions for first two inequalities, respectively.

Proof. Taking $\mu = 1$ in eq. (3.9) yields $|a_3 - a_2^2| \leq \frac{4}{45}$. Using (3.6) and (3.4), we obtain

$$\begin{aligned}
|a_5 - a_3^2| & = \left| \frac{2}{125} \left(c_4 - c_1c_3 - \frac{1}{2}c_2^2 + \frac{3}{4}c_1^2c_2 - \frac{3}{32}c_1^4 \right) - \frac{4}{2025} \left(c_2 - \frac{1}{2}c_1^2 \right)^2 \right| \\
& = \frac{2}{125} \left| \frac{37}{288}c_1^4 + \frac{101}{162}c_2^2 + c_1c_3 - \frac{283}{324}c_1^2c_2 - c_4 \right| \\
& = \frac{2}{125} \left| lc_1^4 + rc_2^2 + 2mc_1c_3 - \frac{3n}{2}c_1^2c_2 - c_4 \right|,
\end{aligned}$$

where $l = \frac{37}{288}$, $r = \frac{101}{162}$, $m = \frac{1}{2}$, $n = \frac{283}{486}$. These values of l, r, m, n satisfy the hypothesis of Lemma 2.4 as it is evident from the facts that $0 < r < 1$, $0 < m < 1$, $4m^2(1-m)^2r(1-r) = 0.0586896$ and

$$8r(1-r)((mn - 2l)^2 + (m(r+m) - n)^2) + m(1-m)(n - 2rm)^2 = 0.00341618.$$

Therefore, by Lemma 2.4,

$$\left| \frac{37}{288}c_1^4 + \frac{101}{162}c_2^2 + c_1c_3 - \frac{283}{324}c_1^2c_2 - c_4 \right| \leq 2.$$

Hence $|a_5 - a_3^2| \leq \frac{4}{125}$. On similar lines, utilizing the inequalities $|a_7| \leq \frac{38}{245}$ and $|a_4| \leq \frac{1}{20}$ as proved in Theorem 3.1, we obtain $|a_7 - a_4^2| \leq |a_7| + |a_4|^2 \leq \frac{3084}{19600}$. \square

We now estimate an upper bound of $|H_{3,1}(f)|$ for $f \in \mathcal{R}_1(\varphi_{3L})$.

Theorem 3.7. If $f \in \mathcal{R}_1(\varphi_{3L})$, then $|H_{3,1}(f)| \leq \frac{481}{90000}$.

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). Then in view of Theorem 3.5 and Theorem 3.6, we have $|a_4 - a_2a_3| \leq \frac{1}{20}$, $|a_3 - a_2^2| \leq \frac{4}{45}$ and $|a_5 - a_3^2| \leq \frac{4}{125}$.

It is found in [24] that

$$|H_{3,1}(f)| \leq |a_5 - a_3^2||a_3 - a_2^2| + |a_4 - a_2a_3|^2.$$

Hence, $|H_{3,1}(f)| \leq \left(\frac{4}{125}\right)\left(\frac{4}{45}\right) + \left(\frac{1}{20}\right)^2 = \frac{481}{90000} \approx 0.00534444$. \square

Theorem 3.8. If $f \in \mathcal{R}_1(\varphi_{3L})$ is given by (1.1), then $|H_{3,2}(f)| \leq \frac{2978957}{1620000000}$.

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). By using (3.3)-(3.6), we obtain

$$\begin{aligned} |a_2a_5 - a_3a_4| &= \frac{1}{11250} \left| 18c_1 \left(c_4 - \frac{11}{36}c_1c_3 \right) - 2c_1^3 \left(c_2 - \frac{23}{32}c_1^2 \right) - \frac{1}{4}c_2(13c_1^3 - 64c_1c_2 + 100c_3) \right| \\ &\leq \frac{1}{11250} \left(18|c_1| \left| c_4 - \frac{11}{36}c_1c_3 \right| + 2|c_1|^3 \left| c_2 - \frac{23}{32}c_1^2 \right| + \frac{1}{4}|c_2| |13c_1^3 - 64c_1c_2 + 100c_3| \right) \\ &\leq \frac{1}{11250} (72 + 32 + 100) \\ &= \frac{34}{1875}. \end{aligned}$$

By using the fact that $|a_2a_5 - a_3a_4| \leq \frac{34}{1875}$ and utilizing the bounds proven in Theorem 3.1, Theorem 3.3 and Theorem 3.4, we obtain

$$\begin{aligned} |H_{3,2}(f)| &\leq |a_6| |a_2a_4 - a_3^2| + |a_5| |a_2a_5 - a_3a_4| + |a_4| |a_3a_5 - a_4^2| \\ &\leq \frac{8}{10125} + \frac{136}{234375} + \frac{3373}{7200000} \\ &= \frac{2978957}{1620000000} \\ &\approx 0.00183886. \end{aligned} \quad \square$$

3.2 Upper Bounds of $|H_{4,1}(f)|$, $|T_4(1)|$, $|T_4(2)|$ for $f \in \mathcal{R}_1(\varphi_{3L})$

Theorem 3.9. If $f \in \mathcal{R}_1(\varphi_{3L})$ is given by (1.1), then $|a_5 - a_2a_4| \leq \frac{4}{125}$. This inequality is sharp.

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). By using (3.6), (3.3) and (3.5),

$$\begin{aligned} |a_5 - a_2a_4| &= \frac{2}{125} \left| \frac{17}{128}c_1^4 + \frac{1}{2}c_2^2 + \frac{37}{32}c_1c_3 - \frac{29}{32}c_1^2c_2 - c_4 \right| \\ &= \frac{2}{125} \left| lc_1^4 + rc_2^2 + 2mc_1c_3 - \frac{3n}{2}c_1^2c_2 - c_4 \right|, \end{aligned}$$

where $l = \frac{17}{128}$, $r = \frac{1}{2}$, $m = \frac{37}{64}$ and $n = \frac{29}{48}$.

The values of l, r, m, n satisfy the hypothesis of Lemma 2.5 and hence

$$|a_5 - a_2 a_4| = \frac{2}{125} \left| l c_1^4 + r c_2^2 + 2m c_1 c_3 - \frac{3n}{2} c_1^2 c_2 - c_4 \right| \leq \frac{4}{125}.$$

The equality hold in $|a_5 - a_2 a_3| \leq \frac{4}{125}$, for $f_4(z) = z + \frac{4}{125} z^5 + \frac{1}{1445} z^{17} \in \mathcal{R}_1(\varphi_{3L})$. \square

Theorem 3.10. If $f \in \mathcal{R}_1(\varphi_{3L})$ is given by (1.1), then $|a_6 - a_3 a_4| \leq \frac{77}{900}$.

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). By using (3.7), (3.5) and (3.4) leads to

$$\begin{aligned} |a_6 - a_3 a_4| &= \frac{1}{900} \left| 10c_5 - 11c_2 c_3 + \frac{17}{4} c_1 c_2^2 + 10c_1 \left(\frac{1}{80} c_1^4 + \frac{17}{40} c_2^2 + \frac{4}{5} c_1 c_3 - \frac{9}{20} c_1^2 c_2 - c_4 \right) \right| \\ &\leq \frac{1}{900} \left(10 \left| c_5 - \frac{27}{40} c_2 c_3 \right| + \frac{17}{4} |c_2| |c_3 - c_1 c_2| + 10 |c_1| \left| l c_1^4 + r c_2^2 + 2m c_1 c_3 - \frac{3n}{2} c_1^2 c_2 - c_4 \right| \right), \end{aligned}$$

where $l = \frac{1}{80}$, $\frac{17}{40}$, $m = \frac{2}{5}$ and $n = \frac{3}{10}$.

These values of l, r, m, n satisfy the hypothesis of Lemma 2.5 and hence

$$\left| \frac{1}{80} c_1^4 + \frac{17}{40} c_2^2 + \frac{4}{5} c_1 c_3 - \frac{9}{20} c_1^2 c_2 - c_4 \right| \leq 2.$$

In view of Lemma 2.4, we have $|c_5 - \frac{27}{40} c_2 c_3| \leq 2$, $|c_3 - c_1 c_2| \leq 2$. Further from Lemma 2.1, we have $|c_1| \leq 2$, $|c_2| \leq 2$. Utilising these inequalities, we obtain that $|a_6 - a_3 a_4| \leq \frac{77}{900}$. \square

Remark 3.1. It is clear from Theorem 3.6 that the Zalcman conjecture is true for $n = 2, 3, 4$ for $f \in \mathcal{R}_1(\varphi_{3L})$. Further, generalized Zalcman conjecture for certain initial values of n, m in view of Theorems 3.5, 3.9, 3.10.

Theorem 3.11. If $f \in \mathcal{R}_1(\varphi_{3L})$, then $|H_{4,1}(f)| \leq 0.00179948585$.

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). It is found in [24] that if $f \in \mathcal{S}$ of the form (1.1), then

$$\begin{aligned} |H_{4,1}(f)| &\leq |a_7 - a_4^2| |H_{3,1}(f)| + |a_6 - a_3 a_4|^2 |a_3 - a_2^2| \\ &\quad + |a_5 - a_2 a_4|^2 |a_5 - a_3^2| + 2|a_6 - a_3 a_4| |a_5 - a_2 a_4| |a_4 - a_2 a_3|. \end{aligned} \quad (3.10)$$

Thus, the required upper bound follows by utilizing the bounds obtained in Theorem 3.2, Theorem 3.5, Theorem 3.6, Theorem 3.7, Theorem 3.9 and Theorem 3.10 in (3.10). \square

Theorem 3.12. If $f \in \mathcal{R}_1(\varphi_{3L})$, then $|T_4(1)| \leq 1.149169$.

Proof. Let $f \in \mathcal{R}_1(\varphi_{3L})$ be given by (1.1). Then by using the bounds of initial coefficients of $f \in \mathcal{R}_1(\varphi_{3L})$ proved in Theorem 3.1, we obtain

$$\begin{aligned} |a_3 - a_2 a_4| &\leq |a_3| + |a_2| |a_4| \leq \frac{4}{45} + \frac{1}{5} \left(\frac{1}{20} \right) = \frac{89}{900}, \\ |a_2 - a_2 a_3| &\leq |a_2| (1 + |a_3|) \leq \frac{1}{5} \left(1 + \frac{4}{45} \right) = \frac{49}{225}, \\ |1 - a_2^2| &\leq 1 + |a_2|^2 \leq 1 + \frac{1}{25} = \frac{26}{25}. \end{aligned}$$

Also, $|a_3 - a_2^2| \leq \frac{4}{45}$, $|a_2a_3 - a_4| \leq \frac{1}{20}$ and $|H_{2,2}(f)| \leq \frac{16}{2025}$ are as proved in Theorems 3.6, 3.5 and 3.3, respectively. In view of [37], we have

$$\begin{aligned} |T_4(1)| &= |(1 - a_2^2)^2 - (a_2a_3 - a_4)^2 + (a_3^2 - a_2a_4)^2 - (a_2 - a_2a_3)^2 + 2(a_2^2 - a_3)(a_3 - a_2a_4)| \\ &\leq |1 - a_2^2|^2 + |a_2a_3 - a_4|^2 + |a_3^2 - a_2a_4|^2 + |a_2 - a_2a_3|^2 + 2|a_2^2 - a_3||a_3 - a_2a_4| \\ &\leq \left(\frac{26}{25}\right)^2 + \frac{1}{400} + \left(\frac{16}{2025}\right)^2 + \left(\frac{49}{225}\right)^2 + 2\left(\frac{4}{45}\right)\left(\frac{89}{900}\right) \\ &\approx 1.149169. \end{aligned}$$

□

Theorem 3.13. If $f \in \mathcal{R}_1(\varphi_{3L})$, then $|T_4(2)| \leq 0.00340790418$.

Proof. Since $f \in \mathcal{R}_1(\varphi_{3L})$, in view of Theorems 3.4, 3.3 and 3.8, we have $|H_{2,3}(f)| \leq \frac{3373}{360000}$, $|H_{2,2}(f)| \leq \frac{16}{2025}$ and $|a_3a_4 - a_2a_5| \leq \frac{34}{1875}$. Further

$$\begin{aligned} |T_2(2)| &= |a_2^2 - a_3^2| \leq |a_2^2| + |a_3^2| \leq \frac{97}{2025}, \\ |a_2a_3 - a_3a_4| &\leq |a_3|(|a_2| + |a_4|) \leq \frac{1}{45}, \\ |a_2a_4 - a_3a_5| &\leq |a_2||a_4| + |a_3||a_5| \leq \frac{1}{100} + \frac{16}{5625} = \frac{289}{22500}. \end{aligned}$$

Following the inequality found in [37], we obtain

$$\begin{aligned} |T_4(2)| &\leq |T_2(2)|^2 + |a_3a_4 - a_2a_5|^2 + |H_{2,3}(f)|^2 + |a_2a_3 - a_3a_4|^2 + 2|H_{2,2}(f)||a_2a_4 - a_3a_5| \\ &\leq \frac{9409}{4100625} + \frac{1156}{3515625} + \frac{11377129}{129600000000} + \frac{1}{2025} + 2\left(\frac{16}{2025}\right)\left(\frac{289}{22500}\right) \\ &= 0.00340790418. \end{aligned}$$

□

4. Concluding Remarks

The upper bounds of Hankel and Toeplitz determinants of order four have been established, shown that the Zalcman conjecture is true for cases $n = 2, 3, 4$. Additionally, the generalized Zalcman conjecture has been validated for initial values of n and m within the class $\mathcal{R}_1(\varphi_{3L})$ in this paper. One can investigate the upper bounds of $|H_{5,1}(f)|$, $|T_5(1)|$ for $\mathcal{R}_1(\varphi_{3L})$ as continuation of the work.

Acknowledgement

The authors sincerely thank the referees for their valuable suggestions, which have significantly improved the quality of the paper.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contributions

All the authors contributed significantly in writing this article. The authors read and approved the final manuscript.

References

- [1] M. F. Ali, D. K. Thomas and A. Vasudevarao, Toeplitz determinants whose elements are the coefficients of analytic and univalent functions, *Bulletin of the Australian Mathematical Society* **97**(2) (2018), 253 – 264, DOI: 10.1017/S0004972717001174.
- [2] M. Arif, L. Rani, M. Raza and P. Zaprawa, Fourth Hankel determinant for the family of functions with bounded turning, *Bulletin of the Korean Mathematical Society* **55**(6) (2018), 1703 – 1711, DOI: 10.4134/BKMS.b170994.
- [3] M. Arif, M. Raza, H. Tang, S. Hussain and H. Khan, Hankel determinant of order three for familiar subsets of analytic functions related with sine function, *Open Mathematics* **17**(1) (2019), 1615 – 1630, DOI: 10.1515/math-2019-0132.
- [4] M. Arif, O. M. Barukab, S. A. Khan and M. Abbas, The sharp bounds of Hankel determinants for the families of three-leaf-type analytic functions, *Fractal and Fractional* **6**(6) (2022), 291, DOI: 10.3390/fractalfract6060291.
- [5] K. O. Babalola, On $H_3(1)$ Hankel determinant for some classes of univalent functions, *Inequality Theory and Applications* **6** (2010), 1 – 7.
- [6] S. Gandhi, Radius estimates for three leaf function and convex combination of starlike functions, in: *Mathematical Analysis I: Approximation Theory* (ICRAPAM 2018), Springer Proceedings in Mathematics & Statistics, Vol. 306, Springer, Singapore (2020), DOI: 10.1007/978-981-15-1153-0-15.
- [7] A. W. Goodman, *Univalent Functions*, Vols. I and II, Polygonal Publishing House, Washinton, New Jersey (1983).
- [8] W. K. Hayman, On the second Hankel determinant of mean univalent functions, *Proceedings of the London Mathematical Society* **s3-18**(1) (1968), 77 – 94. DOI: 10.1112/plms/s3-18.1.77.
- [9] G. Kaur and G. Singh, Fourth Hankel determinant for a new subclass of Bounded turning functions, *Jñānābha* **52**(1) (2022), 229 – 233, DOI: 10.58250/Jnanabha.2022.52129.
- [10] M. G. Khan, B. Ahmad, J. Sokol, Z. Muhammad, W. K. Mashwani, R. Chinram and P. Petchkaew, Coefficient problems in a class of functions with bounded turning associated with sine function, *European Journal of Pure and Applied Mathematics* **14**(1) (2021), 53 – 64, DOI: 10.29020/nybg.ejpam.v14i1.3902.
- [11] M. G. Khan, N. E. Cho, T. G. Shaba, B. Ahmad and W. K. Mashwani, Coefficient functionals for a class of bounded turning functions related to modified sigmoid function, *AIMS Mathematics* **7**(2) (2022), 3133 – 3149, DOI: 10.3934/math.2022173.
- [12] G. Koride, B. S. Rayaprolu, K. Saroja and K. Yakaiah, Fourth Hankel and Toeplitz determinants for a subclass of analytic functions subordinate to $1 + \sin z$, *Mathematica Applicanda (Matematyka Stosowana)* **51**(2) (2023), 321 – 338, DOI: 10.14708/ma.v51i2.7238.
- [13] B. Kowalezyk, A. Lecko and Y. J. Sim, The sharp bound for the Hankel determinant of the third kind for convex functions, *Bulletin of the Australian Mathematical Society* **97**(3) (2018), 435 – 445, DOI: 10.1017/S0004972717001125.
- [14] R. J. Libera and E. J. Ziotkiewicz, Coefficient bounds for the inverse of a function with derivatives in \mathcal{P} , *Proceedings of the American Mathematical Society* **87** (1983), 251 – 257, DOI: 10.1090/S0002-9939-1983-0681830-8.
- [15] W. Ma and D. Minda, A unified treatment of some special classes of univalent functions, in: *Proceedings of the Conference on Complex Analysis*, Nankai Institute of Mathematics, Nankai Univerity, Tianjin, June 19-23, 1992, International Press, pp. 157 – 169 (1994).

- [16] W. Ma, Generalized Zalcman conjecture for starlike and typically real functions, *Journal of Mathematical Analysis and Applications* **234**(1) (1999), 328 – 339, DOI: 10.1006/jmaa.1999.6378.
- [17] S. Mandal, P. P. Roy and M. B. Ahmad, Hankel and Toeplitz determinants of logarithmic coefficients of inverse functions for certain classes of univalent functions, *Iranian Journal of Science* **49** (2025), 243 – 252, DOI: 10.1007/s40995-024-01717-6.
- [18] G. Murugusundaramoorthy, K. Vijaya, K. R. Karthikeyan, S. M. El-Deeb and J.-S. Ro, Bi-univalent functions subordinated to a three leaf function induced by multiplicative calculus, *AIMS Mathematics* **9**(10) (2024), 26983 – 26999, DOI: 10.3934/math.20241313.
- [19] G. Murugusundaramoorthy, M. G. Khan, B. Ahmad, W. K. Mashwani, T. Abdeljawad and Z. Salleh, Coefficient functionals for a class of bounded turning functions connected to three leaf function, *Journal of Mathematics and Computer Science* **28**(3) (2023), 213 – 223, DOI: 10.22436/jmcs.028.03.01.
- [20] C. Pommerenke, On the coefficients and Hankel determinants of Univalent functions, *Journal of the London Mathematical Society* **s1-41**(1) (1966), 111 – 122, DOI: 10.1112/jlms/s1-41.1.111.
- [21] C. Pommerenke, On the Hankel determinants of Univalent functions, *Mathematika* **14**(1) (1967), 108 – 112, DOI: 10.1112/S002557930000807X.
- [22] C. Pommerenke, Univalent functions, in: *Studia Mathematica Mathematiche Lehrbucher*, Vandenhoeck and Ruprecht, Gotingen (1975).
- [23] V. Radhika, S. Sivasubramanian, G. Murugusundaramoorthy and J. M. Jahangiri, Toeplitz matrices whose elements are the coefficients of functions with bounded boundary rotation, *Journal of Complex Analysis* **2016**(1) (2016), 4960704, DOI: 10.1155/2016/4960704.
- [24] S. S. Rao, *A Study on Hankel Toeplitz Determinants and Bilinear Transformations*, Doctoral Thesis, Department of Mathematics, Kakatiya University, Telangana, India (2024), URL: <http://hdl.handle.net/10603/576200>.
- [25] V. Ravichandran and S. Verma, Bound for the fifth coefficient of certain starlike functions, *Comptes Rendus. Mathématique* **353**(6) (2015), 505 – 510, DOI: 10.1016/j.crma.2015.03.003.
- [26] A. Riaz, M. Raza, M. A. Binyamin and A. Saliu, The second and third Hankel determinants for starlike and convex functions associated with three-leaf function, *Heliyon* **9**(1) (2023), e12748, DOI: 10.1016/j.heliyon.2022.e12748.
- [27] K. Sharma, N. K. Jain and V. Ravichandran, Starlike functions associated with a cardioid, *Afrika Matematika* **27** (2016), 923 – 939. DOI: 10.1007/s13370-015-0387-7.
- [28] L. Shi, M. Arif, K. Ullah, N. Alreshidi and M. Shutaywi, On sharp estimate of third Hankel determinant for a subclass of starlike functions, *Fractal and Fractional* **6**(8) (2022), 437, DOI: 10.3390/fractalfract6080437.
- [29] L. Shi, M. Arif, M. Raza and M. Abbas, Hankel determinant containing logarithmic coefficients for bounded turning functions connected to a three-leaf-shaped domain, *Mathematics* **10**(16) (2022), 2924, DOI: 10.3390/math10162924.
- [30] L. Shi, M. G. Khan, B. Ahmad, W. K. Mashwani, P. Agarwal and S. Momani, Certain coefficient estimate problems for three-leaf-type starlike functions, *Fractal and Fractional* **5**(4) (2021), 137, DOI: 10.3390/fractalfract5040137.
- [31] H. M. Srivastava, G. Kaur and G. Singh, Estimates of the fourth Hankel determinant for a class of analytic functions with bounded turnings involving Cardioid domains, *Journal of Nonlinear and Convex Analysis* **22**(3) (2021), 511 – 526.

- [32] S. P. Vijayalakshmi, T. V. Sudharsan and T. Bulboacă, Symmetric Toeplitz determinants for classes defined by post quantum operators subordinated to the limaçon function, *Studia Universitatis Babeş-Bolyai Mathematica* **69**(2) (2024), 299 – 316, DOI: 10.24193/subbm.2024.2.04.
- [33] K. Yakaiah and R. B. Sharma, Fourth Hankel and Toeplitz Determinants for a class of Analytic Univalent Functions subordinated to $\cos z$, *Stochastic Modeling and Applications* **26**(3) (2022), 619 – 623.
- [34] K. Yakaiah, R. B. Sharma, K. Ganesh and K. Saroja, Fourth Hankel and Toeplitz determinants for bounded turning functions subordinate to $1 + \tanh z$, *Advanced Studies: Euro-Tbilisi Mathematical Journal* **16**(3) (2023), 1 – 9, DOI: 10.32513/asetmj/193220082321.
- [35] K. Yakaiah, R. B. Sharma, V. S. Kumar and S. S. Rao, Fourth Hankel and Toeplitz determinants for reciprocal of Bounded turning functions subordinate to $\cos z$, *Communications in Mathematics and Applications* **14**(2) (2023), 969 – 980, DOI: 10.26713/cma.v14i2.2200
- [36] P. Zaprawa, Third Hankel determinants for subclasses of univalent functions, *Mediterranean Journal of Mathematics* **14** (2016), article number 19. DOI: 10.1007/s00009-016-0829-y.
- [37] H.-Y. Zhang and H. Tang, Fourth Toeplitz determinants for starlike functions defined by using the sine function, *Journal of Function Spaces* (2021)(1), 4103772, DOI: 10.1155/2021/4103772.

